

ARTHROPODA.—*Continued.*

(b) Lepidoptera.

Butterflies.—*Papilio macleayanus*, *Delias harpalyce*, *Belenois teutonia*, *Heteronympha merope*, *Epinephile abeona*, *Pyrameis kershawii*, *Pyrameis itea*, *Xenica hobartia*, *Xenica achanta*, *Ialmenus evagores*, *Chrysophanus erinus*, *Lycæna boetica*, *Lycæna Phœbe*, *Taractrocera papyria*, *Hesperia ornata*.

Moths.—*Agarista latinus*, *Agarista lewenii*, *Dasypodia selene-phora*, *Helioches armigera*, *Lucania entranca*, *Camptagramma correlata*, *Camptagramma strangulata*, *Arhodia lutosaria*, *Chærocampa scrofa*, *Procris vividipulvernulenta*, *Eulechria leucopsina*, *Setina trifurcata*, *Rhinodia rostraria*, *Macaria australaria*, *Panagra hypenaria*, *Panagra confluaria*, *Panagra plusiata*, *Coremia vicissata*, *Asthena pulchra*, *Acidalia rubraria*, *Mecyna ornithopteralis*, *Endotricha pyrosalis*, *Thinasotia lativittalis*, *Hednota relatalis*, *Sym-moca herodiella*, *Lœlia australasiæ*, *Nola lugens*, *Philabota fascialis*, *Nyctemera amica*.

VERMES.

- (1) *Turbellaria*.—*Geoplana spenceri*, *dendyi* (sp. n.), *frosti* (sp. n.), *alba*, *mediolineata*, *hoggii*, *munda*, *sulphurea* (now recorded for the first time for Victoria), *macmahoni*, *walhallæ*.
- (2) *Chatopoda*.—Three species, one probably a new species of the genus *Megascolides*.
- (3) *Hirudinea*.—*Geobdella*, sp.
- (4) *Nemertea*.—Two specimens from under the bark of trees.

DESCRIPTION OF A NEW VICTORIAN CONE.

By J. H. GATLIFF.

CONUS SEGRAVEL. Spec. nov.

Spire somewhat concavely elevated, coarsely concentrically striated and slightly grooved, maculated with orange brown; base closely striately grooved, striæ slightly waved. Shoulder angle well defined.

Orange brown, with irregularly-sized tentoriform white reticulations, encircled with many dotted lines of rather darker colour, narrow white band at the shoulder, and another broader one below the middle of the body whorl.

Interior, roseate. Length, 1.25 inches.

Shoreham, Western Port, Victoria.

Differs from the common species, *C. aemone*, Lam., owing to the absence of the ridged striæ on the body whorl, so strongly characteristic of that species. Mr. Segrave (after whom the shell is named) found four other specimens, much worn.

It belongs to Tryon's Section 7, Ammirales, of the genus *Conus*.

Two views of the shell are given in the accompanying plate.